# Latest Triassic conodonts of the Slovenian Basin and some remarks on their evolution

# Zgornjetriasni konodonti Slovenskega bazena in njihov evolucijski razvoj

Tea KOLAR-JURKOVŠEK

Geološki zavod Slovenije, Dimičeva ulica 14, SI-1000 Ljubljana; e-mail: tea.kolar@geo-zs.si

Prejeto / Received 7. 4. 2011; Sprejeto / Accepted 5. 5. 2011

*Key words:* conodonts, Late Triassic, biostratigraphy, Mt. Kobla, Slovenian Basin *Ključne besede:* konodonti, zgornji trias, biostratigrafija, Kobla, Slovenski bazen

#### Abstract

A stratigraphical importance of some latest Triassic conodont taxa from the Slovenian Basin with special regard on their distribution in the Slatnik Formation of the Mt. Kobla section (Julian Alps) is presented. Description of a new species *Misikella buseri* n. sp. is given. It is marked by a reduced segminate element and it represents a distinct stage at the decline of *Misikella* evolution.

#### Izvleček

Prikazan je stratigrafski pomen nekaterih pomembnih zgornjetriasnih konodontih rodov v Slovenskem bazenu s sposebnim poudarkom na njihovem pojavljanju v Slatenski formaciji profila Kobla (Julijske Alpe). Opisana je nova vrsta *Misikella buseri* n. sp., za katero je značilen zmanjšan segminatni element in predstavlja eno od zaključnih stopenj ob koncu razvoja rodu *Misikella*.

# Introduction

The phylum Conodonta was one of the groups that became extinct at the Triassic-Jurassic boundary (TJB) when one of the five largest mass extinctions of the Phanerozoic marine record occurred (SEPKOSKI & RAUP, 1986). There are very few reports of Jurassic survivals and Kozur (1993) documented Neohindeodella detrei from the earliest Hettangian of Csővár in Hungary. During the latest Triassic conodonts therefore suffered major decline in diversity and their extinction can be interpreted as the cummulative result of several causal factors and not of a single catastrophic event (CLARK, 1983). There is an evident decline of conodont taxa from Carnian-Norian boundary strata. Many gondolellids last appeared in the Norian, but a few range into the lower Rhaetian, for example *E*. *bidentata* as the last epigondolellid representative. Only few gondolellids, such as Norigondolella and the platform-less Parvigondolella and Misikella, persisted during the latest Triassic.

The study of conodonts has resulted in the recognition of several stratigraphically significant conodont species belonging to the genera: *Budurovignathus, Epigondolella, Gladigondolella,*  Metapolygathus, Misikella, Neogondolella, Nicoraella, Norigondolella and Paragondolella in the Middle and Late Triassic stata of the Slovenian Basin (Buser et al., 2007, 2008). The youngest Triassic conodont assemblages are marked by the presence of *Misikella*, a genus already described from western Slovenia of the Pokljuka plateau and from the Mt. Šija area (Kolar-Jurkovšek et al., 1983; Kolar-Jurkovšek, 1994).

The aims of this paper are to give a brief review of the Upper Triassic conodont genera known from Slovenia (Kolar-Jurkovšek, 1991; Buser et al., 2007, 2008), and to describe a new *Misikella* species, *Misikella buseri* n. sp., from the Rhaetian strata of the Slovenian Basin, sampled in the Mt. Kobla section (Fig. 1).

## **Previous research**

The existence of Late Triassic deep-water sediments in western Slovenia was recognized by many authors (WINKLER, 1923; AUBOUIN, 1960; COUSIN, 1973). They were later object of several basic studies conducted by Buser (BUSER, 1986, 1987, 1989, 1996, 2003; BUSER & DEBELJAK, 1996; BUSER et al., 2007, 2008; BUSER & OGORELEC, 2008)



Fig. 1. Actual position of geotectonic units in western Slovenia with extension of sediments of the ancient Julian and Dinaric Carbonate Platforms and the intermediate Slovenian Basin with the position of the Mt. Kobla section (star). Modified by BUSER et al. (2007).

and are currently object of multidisciplinary study carried out by Rožič and co-workers (Rožič, 2008, 2009; Rožič & Kolar-Jurkovšek, 2007; Rožič et al., 2009; Gale, 2010).

The Slovenian Basin extends in an east-west direction of central Slovenia and became established during the Ladinian following disintegration of the Slovenian Carbonate Platform and it persisted continuously until the Late Cretaceous (BUSER, 1989; BUSER et al., 2007, 2008). The basin pinched out in the Soča river valley and was not connected to the Belluno Basin (BUSER, 1986). From the eastern part of present Slovenia, it continues across Croatia north of Zagrebačka gora to Hungary. The present exposed parts of the Slovenian Basin measure in length about 170 km and in width about 40 km (BUSER et al., 2007, 2008).

After the basic research by BUSER (1986, 1987; BUSER & OGORELEC, 2008) the Mt. Kobla section was sampled and measured also by Rožič et al. (2009). They discriminated the non-dolomitized uppermost part of the Bača Dolomite Formation by formalizing this unit and introducing the Slatnik Formation formed of hemipelagic limestone alternating with resedimented limestone indicating a progradation of sedimentary environments from the the basin plain to the lower slope (Rožič, 2008). The Slatnik Formation is preserved in the northern part of the basin where the latest Triassic succession experienced less intense diagenetic alteration. The Late Triassic part of the Mt. Kobla section is documented by three upward prograding high-frequency cycles. The Triassic part of the section is followed by the Krikov Formation of Jurassic age. The authors also provided conodont data and they recognized three latest Triassic conodont zones: Epigondolella bidentata, Parvigondolella andrusovi-Misikella hernsteini and Misikella hernsteini-Misikella posthernsteini Zones (Rožič et al., 2009). A more detailed sampling of the interval with the finding of M. posthernsteini provided new data and enabled a more precise documentation of the latest zone. Thus, the Misikella posthernsteini Assemblage Zone can be divided into two units: the Misikella hernsteini-Misikella posthernsteini and the Misikella koessenensis Subzones (GALE et al., submitted). The recovered conodont faunas include representatives of Epigondolella, Misikella, Norigondolella, Oncodella, Parvigondolella and Zieglerioconus. The new species of Misikella was collected in the interval from 76,0 to 77,8 m of the Mt. Kobla section (Fig. 2). A detail description of the investigated section along with biostratigraphic conclusions based on conodonts and foraminifers is given in GALE et al. (submitted).



Fig. 2. Geological column of the Mt. Kobla section with conodont distribution. Modified after GALE et al. (submitted).

## **Geological setting**

The Mt. Kobla section (x =5121,590, y = 5420,550, z =1498 m) is located on the southern brim of the the Julian Alps (Fig. 1, 2) and preserves the Upper Norian - Lower Jurassic succession of the Slovenian Basin (Rožič et al., 2009). This area is part of the Tolmin Nappe that together with the overlying Julian Nappe structurally forms the eastern part of the Southern Alps (PLACER, 1999, 2008). The Slovenian Basin is one of the three paleogeographic units that existed during the Late Triassic in the territory of Slovenia. These are (from north to south): the Julian Carbonate Platform, the Slovenian Basin and the Dinaric Carbonate Platform (BUSER, 1986, 1989, 1996) (Fig. 1).

IAN

SEVATIAN

# Significant conodont taxa in the latest Triassic

Norigondolella is represented by the long-ranging species N. steinbergensis. It is an extremely facies dependant species that can be found in fully pelagic sediments of open sea (cherty limestone) of the latest Triassic and therefore it is not stratigraphically important outside this facies (Kozur & Mock, 1991). In the Mt. Kobla section in Slovenia, the LAD of N. steinbergensis is documented in the upper Misikella posthernsteini A. Z. (= Misikella koessenensis - Misikella posthernsteini Subzone) at 88,7 m.

Genus Parvigondolella evolved from Epigondolella bidentata as the ultimate stage in epigondolellid evolution. The genus however is quite rare as can be explained by its existence at the end of evolution of an important stock and it is represented by only few species: P. andrusovi, P. ? lata, P. rhaetica, P. vrielyncki. Some authors regard the platform-less gondolellid forms as juvenile forms of E. bidentata named as the andrusovi stage (KRYSTYN, 1980; KRYSTYN et al., 2007). However, such a view was rejected based on different stratigraphic occurrences, as Parvigondolella ranges up high in the Rhaetian interval that is beyond the highest occurrence of E. bidentata or any other platform-bearing metapolygnathid (Kozur & Mock, 1991). The latter authors also supported their statement by noting the size of the different stages and that certain parameters (size and number of denticles, height of blade) of younger onotogenetic stages cannot exceed the parameters of the largest adult specimens. KOZUR & MOCK (1991) documented their view based on reports of the quite widespread appearance of Parvigondolella in the Rhaetian of Alps, Carpathians, Hungary, Lagonegro Basin that is also confirmed now with data from Slovenia.

Some ramiform elements are also present but are rather rare. Nevertheless in a few cases they permit apparatus reconstruction. Among them Kozur & MOSTLER (1991) reported also a stratigraphically most important species *Neohindeodella detrei* as an index species for the youngest strata above the last occurrence of *Misikella* and these authors stated the form seems to be homeomorph with forms known from older strata of the Late Triassic.

Zieglerioconus has been known to appear with only one representative, Z. rhaeticus, but new Norian species have yet not been described (Channell et al., 2003). It is a secondary single cone conodont with a widely expanded and excavated basal cavity. Zieglerioconus was first described from the Rhaetian (M. koessenensis Subzone) of Csővár in Hungary, and it is rare species that has been so far reported only from Great Britain what means it appears in sediments of the pelagic Tethys and non-pelagic of outer Tethys in the Germanic Basin (Kozur & Mock, 1991; Swift, 1989; Pálfy et al., 2007). The hitherto known occurrences of the species are quite typical for the M. koessenensis Subzone, but Zieglericonus begins in the uppermost Carnian and occurs throughout the Norian (CHANNELL et al., 2003; KOZUR, pers. comm.). New data from Slovenia confirm the existence of *Zieglerioconus* in the late Sevatian. In the Kobla section. Zieglericonus ranges from Parvigondolella andrusovi-Misikella hernsteini A. Z. to lower Misikella koessenensis Sz. of the Misikella posthernsteini Zone. The origin of Zieglerioconus is still vet not known, but Kozur & Mock (1991, p. 277) suggested its forerunner could be a Misikella of the M. longidentata group or Oncodella paucidentata. According to recent data from Slovenia, Zieglerioconus and M. hernsteini made their first appearance together in the Misikella hernsteini-Parvigondolella andrusovi A. Z. in absence of Oncodella and thus, documenting the existence of Zieglerioconus in the latest Norian strata (Fig. 2; Pl. 1, Fig. 2). The forerunner of Zieglerioconus and M. hernsteini is therefore supported to be linked with the M. longidentata group.

**Oncodella paucidentata** is a characteristic element that already appeared in the uppermost Sevatian and it is also a facies controlled species (KOZUR & MOCK, 1991). In Slovenia, it ranges from the upper part of the *Parvigondolella andrusovi-Misikella hernsteini* A.Z. up to an un-named *Misikella* Zone that corresponds to the known range of the species elsewhere.

Genus Misikella is most important for the stratigraphy of the Late Sevatian-Rhaetian interval. Some biostratigraphic schemes have been reported so far and they are slightly different (MOSTLER et al., 1978; GAZDZICKI et al., 1979; KRYSTYN, 1980, 1987; KOZUR, 2003; KRYSTYN et al., 2007). In 1991, KOZUR and MOCK made a report on stratigraphic value of certain conodont taxa. The main intent was to eliminate *M. rhaetica* (it appeared already in the uppermost Sevatian) as an index form for the zone introduced by MOSTLER et al. (1978) as it is strongly facies controlled and occurs already at levels with frequent M. hernsteini. From the upper part of the M. posthernsteini A.Z. they gave a description of the Misikella koessenensis Subzone (originally described by Kozur 1989) as a substitute of the M. rhaetica Zone of earlier schemes, and the author noted also to a shorter range of Misikella koessenensis. The first appearance of Misikella posthernsteini is an important biomarker used to define the Norian-Rhaetian boundary (Kozur & Mock, 1974b, 1991; Kozur, 1996; KRYSTYN, 1980, 1987; BIRKENMAJER et al., 1990). On the other hand, M. ultima is a facies independant species and therefore it is very good boundary marker of the M. ultima A. Z. (the lower boundary is defined by FAD of the index species, but the upper boundary is marked by the dissapearance of *Parvigondolella* and *Misikella*) (Kozur & Mock, 1991).

#### **Evolution of Misikella**

The evolutionary trend M. hernsteini – M. posthernsteini – M. ultima is well known (Kozur & MOCK, 1991). The phylomorphogenetic line between M. hernsteini and M. posthernsteini was documented first by MOSTLER et al. (1978) and was recently studied by GIORDANO et al. (2010) in which transitional forms are arranged in three evolutionary steps. Misikella hernsteini is characterized by a long blade composed of 5-6 denticles and an appropriatelly long and narrow basal dropshaped basal cavity. The step 1 in the evolution of M. hernsteini – M. posthernsteini is marked by a shorter blade with a decreased number of denticles (four in number); the basal cavity is enlarged and has a markedly expanded posterior part with a rounded or triangle-like outline but is devoid of any incision. The increase of the height of denticles is obvious. The length : height ratio (L : H) in M. hernsteini is 2 : 1,5 whereas in the step 1 of the H. hernsteini – H. posthernsteini evolutionary trend is almost 1:1. This step corresponds description of *H. koessenensis*. In general, there are very rare *Misikella* specimens bearing denticles behind the cusp and all of them belong to *M. rhaetica*. The size of *M. koessenensis* is comparable to the size of *M. hernsteini*, whereas in the *M. posthernsteini* an increase in height may be seen. In this study, short and high elements that reveal no posterior denticle, composed of 3-5 denticles and an appropriate ratio L : H are included to M. koessenensis sensu lato rather than make a compromise to regard these forms as Misikella n. sp. B, as suggested by Kozur (pers. comm.). Mostler originally wrote that most specimens of *M. koessenensis* have slender denticle behind the cusp. Thus, the Slovenian specimens fit the original diagnosis of the taxon. However, most conodont workers so far included only specimens with posterior denticle(s) to this species, and therefore an emendation of *M. koessenensis* should be done. In the faunas of the investigated Kobla section such elements make their first apperance at 58 m, 3 metres above the first occurrence of M. hernsteini. The following two evolutionary steps 2 and 3 leading towards M. posthernsteini are marked by a progressive degree of incision in the posterior part of the basal cavity that causes a heart-shaped lower outline, and parallel to this also a reduction of denticles ending in three-dentate elements with a noticeable larger size.

Two lines can be seen in evolution of *Misikella*. They are reprepresented by *M. hernsteini* and *M. posthernsteini*. The *M. hernsteini* evolutionary line is marked by a slight reduction of overall unit size, whereas in the *M. posthernsteini* line an enlargement of overall size is evident. Both lines are marked by a reduction in the number of denticles. The evolutionary trend in the two lines is practically identical if one takes into consideration the outline of the lower side that is drop-shaped in *M. hernsteini* and heart-shaped in *M. posthernsteini*, respectivelly.

We agree with observation of GIORDANO et al. (2010) that the outline of the basal side is important and it is used as the main criterion to distinguish M. hernsteini and M. posthernsteini. Their evolution from units with an oval basal area in *M. hernsteini* is documented with a series of transitional forms leading to elements with an evident inflexion ending in a heart-shaped basal side of M. posthernsteini. Arguments of Moix et al. (2007) claiming that the lower area is important for recognition of Sevatian-Rhaetian taxa is in general supported. They argued the lower side of *M. rhaetica* does not extend to the full length of the lower side of the element but extends only 2/3. Therefore they assigned »M.« rhaetica to Parvigondolella. The original diagnosis of M. rhaetica includes elements bearing 1-3 smaller denticles behind the cusp and the basal cavity extends only 2/3 of the lower side in most specimens (MOSTLER et al. 1978). In this study, the elements of Misikella with 1-2 denticles of smaller size behind the cusp and with a widely opened

basal area extending nearly the full length are attributed to *M. rhaetica* as illustrated in Rožič et al. (2009, Fig. 9e). Thus, the species name *Misikella rhaetica* sensu lato is used herein. Similar specimens are regarded as long *M. hernsteini* by Kozur (pers. comm.) or are assigned to *Misikella* n. sp. A by MUTTONI et al. (2010). Emendation of *M. rhaetica* is therefore needed as the original diagnosis includes also specimens with longer basal area. However, based on the material from Slovenia, we may conclude that there exists a strong homeomorphism in morphology of *Misikella rhaetica* and *Parvigondolella rhaetica*.

The known origin of *Misikella* started with *M. hernsteini*, as its forrunner is yet still not known. However, it could be linked with *M. lon-gidentata*. *M. hernsteini* appeared during the late Sevatian in the *M. hernsteini* – *P. andrusovi* A. Z. Soon after appearance of *M. hernsteini* two lines of evolution can be noted that markedly differ in outline of the basal area. One line retained the drop-shaped basal area, whereas a heart-shaped basal area is evident in the second:

- Evolutionary line: M. hernsteini (M. rhaetica) M. koessenensis M. buseri. Already in the lower M. hernsteini P. andrusovi A. Z. two species, M. koessenensis and M. rhaetica evolved. According to data from Slovenia the two taxa appeared simultaneously where specimens of M. koessenensis with no denticles posterior to the cusp were collected. Such forms gave rise to M. buseri that represents a final step in this evolutionary line marked by obvious reduction of the overall size, decrease in number of denticles and proportional increase of the height of denticles.
- Evolutionary line: M. hernsteini M. posthernsteini M. ultima M. kovacsi: This evolutionary line is marked by the evolution of a heart-shaped basal area in M. posthernsteini that was followed by M. ultima and M. kovasci. Misikella posthernsteini evolved from M. hernsteini through several transitional forms and marks the lower bounday of the M. posthernsteini A. Z. The evolutionary trend M. posthernsteini – M. ultima – M. kovasci is characterized by a decrease of overall size and in a marked reduction of number of denticles.

The two phylogenetic lineages share similar stages of evolution. *M. koessenensis* and *M. posthernsteini* evolved after common ancestor of both lines. At this stage two lines with a distinct forming of the basal area are evident. According to the Slovenian data, *M. koessenensis* and *M. rhaetica* evolved simultaneously from *M. hernsteini*. The intermediate stage is represented by *M. rhaetica* and *M. ultima*. The two species have been so far reported as a paralel step in *Misikella* evolution as they both reveal a secondary bar, but they





Fig. 3. Possible evolution of *Zieglerioconus* and *Misikella* from an unknown origin in the Middle Norian. The chart is based on occurrences in the Slovenian Basin except for the *M. ultima* and *M. kovacsi* occurrences are after KOZUR & MOCK (1991) and PÁLFY et al. (2007); conodont zonation after KOZUR & MOCK (1991). Abbreviations: *M. – Misikella*, *O. – Oncodella*, *Z. – Zieglerioconus*.

retained the similar outline of its ancestor. Based on available data it seems that *M. rhaetica* was a parallel step in *Misikella* evolution. The final step in evolution of both lines is represented by *M. buseri* and *M. kovacsi* with forms characterized by very reduced elements bearing only two denticles. As the following expressed trend in evolution of *Misikella* its ultimate stage could be expected in secondary single cone forms, similar to *Zieglerioconus*. However, their direct relationship is excluded due to their co-occurrences in the Slovenian section. Available data suggest the origin of *Zieglerioconus* and *Misikella* from a common and still unknown ancestor (Fig. 3).

The evolution of the last representatives of gondolellid stock in the latest Triassic (platform type *Epigondolella* and non-platform types Misikella and Parvigondolella) share certain similarities. The evolutionary steps of Epigondolella have already been studied by TRAMMER (1974) who documented a trend in reduction of element size, reduction of platform length, decrease in number of lateral denticles along with increasing height of denticles in the carina. A strong reduction in size and total reduction of the platform was shown in the Middle Triassic lineage Neogondolella – Celsigondolella by Kozur (1968). The end form, Celsigondolella watznaueri is very similar to Parvigondolella. A similar evolutionary trend can be observed also in the non-platform types, Misikella and Parvigondolella. Our study documents that a strong homeomorphism occurred in the evolution of the three genera leading to very reduced forms. This phenomenon was observed in the evolution of certain Late Triassic genera as has already been pointed out by Kozur & MOCK (1974a). The specimen here illustrated (Pl. 1, Fig. 1) reveals a narrow basal area that is not typical for representatives of the genus *Misikella* at this level and morphologically stands close to Parvigondolella, therefore this element is here assigned to the genus »Misikella«.

# Systematic paleontology

# Genus Misikella Kozur & Mock 1974

## Type species: Spathognathodus hernsteini Mostler 1967

## Misikella buseri n. sp. Pl. 1, Fig. 3

Derivatio nominis. In honour of Prof. Dr. Stanko Buser, a Slovenian geologist for his outstanding contribution to the knowledge of the Slovenian Basin and the Slovenian geology in general.

Holotypus. The specimen on Pl. 1, Fig. 3, sample K2/76,0, repository number GeoZS 4242.

Locus typicus. The Kobla section, Tolmin Nappe, western Slovenia.

Stratum typicum. Sample K2/ 76,0, Rhaetian, lower *Misikella posthernsteini* A. Z. (= *Misikella hernsteini* – *Misikella posthernsteini* Subzone), Slatnik Formation.

Material. 5 specimens.

Diagnosis. The P1 element is a very reduced segminate element comparable to other species in *Misikella* marked by a big cusp and one more small denticle, a basal cavity rounded posteriorly and devoid of any incision.

Description. This species is characterized by very reduced segminate element with a large cusp and one more small denticle: in the holotype the denticle next to the cusp is very reduced. The basal area is very opened and has an oval posterior edge. The basal cavity is widely expanded appropriatelly to the large size of the cup.

Occurrence. Late Triassic: early Rhaetian, lower *Misikella posthernsteini* A. Z. (= *Misikella hernsteini* – *Misikella posthernsteini* Subzone), Slatnik Formation.





Conodonts from the Mt. Kobla section, Julian Alps, Slovenia; Slatnik Formation; Figs. 1, 2 – scale bar 100  $\mu$ m, Fig. 3 – scale bar 50  $\mu$ m.

- 1 "Misikella" sp., Pa element, sample K2/64,0 (GeoZS 4234); Parvigondolella andrusovi-Misikella hernsteini A. Z.
- 2 Zieglerioconus rhaeticus Kozur & Mock, Pa element, sample K2/58,0 (GeoZS 4228); Parvigondolella andrusovi-Misikella hernsteini A. Z.
- 3 Misikella buseri n.sp., Pa element, holotype, sample K2/76,0 (GeoZS 4242); lower Misikella posthernsteini A. Z. (= Misikella hernsteini – Misikella posthernsteini Subzone).
- a lateral view, b oblique lower view, c lower view.

The illustrated conodont specimens were photographed at the JEOL JSM 6490LV Scanning Electron Microscope at the Geological Survey of Slovenia / Geološki zavod Slovenije and are housed in the micropaleontological collection of the same institution.

#### Acknowledgements

The author is indebted to Merlynd Nestell (Arlington, Texas) for editing the manuscript and wishes to extend her thanks to Heinz Kozur (Budapest, Hungary) for helpful suggestions and communications on conodonts. Milan Sudar (Belgrade, Serbia) and Bojan Ogorelec (Ljubljana, Slovenia) provided constructive reviews of the manuscript. Facilities and technical staff of the Geological Survey of Slovenia are gratefully acknowledged. The investigation was financially supported by the Slovenian Research Agency (programme number P1-0011).

#### References

- AUBOUIN, J. 1960: Essai sur l'ensemble italo-dinarique et ses rapports avec l'arc alpin. Bull. Soc. Geol. France, 7/2: 487–526.
- BIRKENMAJER, K., KOZUR, H. & MOCK, R. 1990: Exotic Triassic limestone pebbles from the Pieniny Klippen Belt of Poland. Ann. Soc. Geol. Polon., 60: 3–44.
- BUSER, S. 1986: Explanatory book, Sheet Tolmin and Videm (Udine) L33-64, L33-63. Basic geological map of SFRY 1 : 100,000. Zvezni geološki zavod, Beograd.
- BUSER, S. 1987: Osnovna geološka karta SFRJ 1 : 100 000, list Tolmin. Zvezni geološki zavod, Beograd.
- BUSER, S. 1989: Development of the Dinaric and the Julian carbonate platforms and of the intermediate Slovenian Basin (NW Yugoslavia). Boll. Soc. Geol. It., 40: 313–320.
- BUSER, S. 1996: Geology of Western Slovenia and its paleogeographic evolution. In: DROBNE, K., GORIČAN, Š. & KOTNIK, B. (Eds.): The role of impact processes and biological evolution of Planet Earth: International workshop, Postojna, 27. 9. – 2. 10. 1996, ZRC SAZU, Ljubljana, 111– 123.
- BUSER, S. 2003: Géologie de la Slovénie occidentale. In: DROBNE, K., PUGLIESE, N. & TAMBAREAU, Y. (Eds.): De la mer Adriatique aux Alpes Juliennes (Italie nord-orient et Slovénie occidentale) – un percours géologique sans frontieres. Ljubljana: Znanstveno raziskovalni center SAZU; Trieste: Dipartimento di scienze geologiche, ambientali e marine, Universitá, 27–31.
- BUSER, S. & DEBELJAK, I. 1996: Lower Jurassic beds with bivalves in south Slovenia. Geologija, 37/38 (1994/95): 23–62.
- BUSER, S., KOLAR-JURKOVŠEK, T. & JURKOVŠEK, B. 2007: Triasni konodonti slovenskega bazena = Triassic conodonts of the Slovenian Basin. Geologija, 50/1: 19–28.
- BUSER, S., KOLAR-JURKOVŠEK, T. & JURKOVŠEK, B. 2008: The Slovenian Basin during the Triassic in the Light of Conodont Data. Boll. Soc. Geol. It. (Ital. J. Geosci.), 127/2: 257–263.
- BUSER, S. & OGORELEC, B. 2008: Globljevodne triasne in jurske plasti na Kobli = Deep-water Triassic and Jurassic beds from Mt. Kobla, W Slovenia. Geologija, 51/2: 181–189.
- CHANNELL, J. E. T., KOZUR, H. W., SIEVERS, T., MOCK, R., AUBRECHT R. & SYKORA, M. 2003: Carnian -

Norian biomagnetostratigraphy at Silická Brezová (Slovakia): correlation to other Tethyan sections and to the Newark Basin. Palaeogeogr., Palaeoclimatol., Palaeoecol., 191: 65–109.

- Cousin, M. 1973: Le sillon slovène: les formations triasiques, jurassiques et neocomiennes au Nord-Est de Tolmin. Bull. Soc. geol. Fr., 7/15: 326–339.
- CLARK, D. L. 1983: Extinction of conodonts. J. Paleont., 57: 652–661.
- GALE, L. 2010: Microfacies analysis of the Upper Triassic (Norian) "Bača Dolomite": early evolution of the western Slovenian Basin (eastern Southern Alps, western Slovenia). Geol. Carpath., 61: 293–308.
- GALE, L., KOLAR-JURKOVŠEK, T., ŠMUC, A. & ROŽIČ, B. (submitted): Integrated foraminiferal and conodont biostratigraphy from the Rhaetian strata of the Slovenian Basin: new data from the Mt Kobla section (eastern Southern Alps, western Slovenia).
- GAźDZICKI, A., KOZUR, H. & MOCK, R. 1979: The Norian-Rhaetian boundary in the light of micropaleontological data. Geologija, 22: 71–112.
- GIORDANO, N., RIGO, M., CIARAPICA, G., BERTINELLI, A. 2010: New biostratigraphical constraints for the Norian/Rhaetian boundary: data from Lagonegro Basin, Southern Apennines, Italy. Lethaia, 43/4: 573-586,

doi:10.1111/j.1502-3931.2010.00219.x

- Kolar-Jurkovšek, T. 1994: Mikrofauna from the Upper Triassic of Karavanke Mts. (Slovenia). Mém. de Géol., 22, Proceedings of the Triassic Symposium, Lausanne, 20-25 October 1991: 53–62.
- KOLAR-JURKOVŠEK, T. 1991: Mikrofavna srednjega in zgornjega triasa Slovenije in njen biostratigrafski pomen. Geologija, 33: 21–170.
- KOLAR-JURKOVŠEK, T., BUSER, S. & JURKOVŠEK, B. 1983: Zgornjetriasne plasti zahodnega dela Pokljuke. Rud.-metal. zb., 30/2-3: 151–185.
- KOZUR, H. 1968: Conodonten aus dem Muschelkalk des germanischen Binnenbeckens und ihr stratigraphischer Wert. Teil I: Conodonten vom Platformtyp und stratigraphische Bedeutung der Conodonten aus dem oberen Muschelkalk. Geologie, 17/8: 930–946.
- KOZUR, H. 1989: Significance of Events in Conodont Evolution for the Permian and Triassic Stratigraphy. Courier Forsch., Inst. Senckenberg, 117: 385–408.
- KOZUR, H. 1993: First evidence of Liassic in the vicinity of Csövár (Hungary) and its paleogeographic and paleotectonic significance. Jb. Geol. B.-A., 136/1: 89–98.
- KOZUR, H. 1996: The Conodonts Hindeodus, Isarcicella and Sweetohindeodus in the Uppermost Permian and Lowermost Triassic. Geol. Croat., 49/1: 81–115.
- KOZUR, H. 2003. Integrated ammonoid-, conodont and radiolarian zonation of the Triassic. Hallesches Jahrb. Geowiss., 25: 49–79.
- KOZUR, H. & MOCK, R. 1974a: Zwei neue Conodonten-Arten aus der Trias des Slowakischen Karstes. Čas. Min. Geol., 19: 135–139.

- KOZUR, H. & MOCK, R. 1974b: Misikella posthernsteini n.sp., die jüngste Conodontenart der tethyalen Trias. Čas. Min. Geol., 19: 245–250.
- KOZUR, H. & MOCK, R. 1991: New Middle Carnian and Rhaetian conodonts from Hungary and the Alps. Stratigraphic importance and tectonic implications for the Buda Mountains and adjacent areas. Jb. Geol. B.-A., 134: 271–297.
- KRYSTYN, L. 1980: Triassic conodont localities of the Salzkammergut region (Northern Calcareous Alps). In: SCHÖNLAUB, H.P. (Ed.): Second European conodont symposium – ECOS II, Guidebook, Abstracts. Abh. Geol. B.-A., 35: 61–98.
- KRYSTYN, L. 1987: Zur RHÄT-Stratigraphie in den Zlambach-Schichten (vorläufiger Bericht). Sitz. Ber. Akad. Wiss. Wien, math.-nat. Kl., 196: 21–36.
- KRYSTYN, L., BOUQUEREL, H., KUERSCHNER, W., RI-CHOZ, S. & GALLET, Y. 2007: Proposal for an austrian candidate GSSP for the base of the Rhaetian stage. In: LUCAS, S. G. & SPIELMANN, J. A. (Eds.): The Global Triassic. New Mexico Museum. Nat. Hist. & Sc. Bull., 41: 189–199.
- MOIX, P., KOZUR, H. W., STAMPFLI, G. M. & MOSTLER, H. 2007: New paleontological, biostratigraphic and paleogeographic results from the Triassic of the Mersin Mélange, SE Turkey. In: LUCAS, S. G. & SPIELMANN, J. A. (Eds.): The Global Triassic. New Mexico Mus. Nat. Hist. Sci. Bull., 41: 282–311.
- MOSTLER, H. 1967: Conodonten und Holothuriensklerite aus den norischen Hallstätterkalken von Hernstein (Niederösterreich). Verh. Geol. Bundestanst., 1–2: 177–188.
- Mostler, H., Scheuring, B. & Ulrichs, M. 1978: Zur Mega- und Mikrofauna und Mikroflora der Kössenerschichten (alpine Obertrias) wom Weissloferbach in Tirol unter besondere Berücksichtigung der in der suessi- und marshi- Zone auftretenden Conodonten. Schrift. Erdwiss. Komm. Österr. Akad. Wiss., 4: 141–174.
- MUTTONI, G., KENT, D. V., JADOUL, F., OLSEN, P. E., RIGO, M., GALLI, M. T., NICORA, A. 2010 : Rhaetian magneto-biostratigraphy from the Southern Alps (Italy): constraints on Triassic chrono-

logy. Palaeogeogr., Palaeoclimatol., Palaeoecol., 285: 1–16.

- PÁLFY, J., DEMÉNY, A., HAAS, J., CARTER, E. S., GÖRÖG, Á., HALÁSZ, D., ORAVECZ-SCHEFFER, A., HETÉNYI, M., MÁRTON, E., ORCHARD, M. J., OZS-VÁRT, P., VETŐ, I. & ZAJZON, N. 2007: Triassic-Jurassic boundary events inferred from integrated stratigraphy of the Csővár section, Hungary. Palaeogeogr., Palaeoclimatol., Palaeoecol., 244: 11–33.
- PLACER, L. 1999 : Contribution to the macrotectonic subdivision of the border region between Southern Alps and External Dinarides. Geologija, 41 (1998): 223–255.
- PLACER, L. 2008: Principles of the tectonic subdivision of Slovenia. Geologija, 51/2: 205–217.
- Rožič, B. 2008: Upper Triassic and Lower Jurassic limenstones from Mt Kobla in the northern Tolmin Basin: tectonically repeated or continous succession? RMZ-Materials and Geoenvironment, 55/3: 345–362.
- Rožič, B. 2009: Perbla and Tolmin formations: revised Toarcian to Tithonian stratigraphy of the Tolmin Basin (NW Slovenia) and regional correlations. Bull. Soc. Geol. France, 180/5: 409–423.
- Rožič, B. & Kolar-Jurkovšek, T. 2007: Zgornjetriasni apnenčevi razvoji Slovenskega bazena na Kobli in Slatniku. Geološki zbornik, 19: 96–99.
- Rožič, B., Kolar-Jurkovšek, T. & Šmuc, A. 2009: Late Triassic sedimentary evolution of Slovenian Basin (eastern Southern Alps): description and correlation of the Slatnik Formation. Facies, 55/1: 137–155.
- SWIFT, A. 1989: First records of conodonts from the Late Triassic of Britain. Palaeontology, 32: 325–333.
- SEPKOSKI, J. J. & RAUP, D.M. 1986: Periodicity in marine extinction events. In: Elliot D.K. (Ed.): Dynamics of Extinction. Wiley, New York: 3–36.
- TRAMMER, J. 1974: Evolutionary Trends and Pattern of Extinction of Triassic Conodonts. Acta Pal. Polonica, 19/2: 251–264.
- WINKLER, A. 1923: Über den Bau der östlichen Südalpen. Mitt. Öster. Geol. Gesellschaft, 16: 1–272.